

The 2024 Health Research and Policy Conference 20-21 May 2024

RWANDA BIOMEDICAL CENTRE, KIGALI, RWANDA

Theme: Advancing Research for a Healthy Rwanda

The 2024 Health Research and Policy Conference, held on May 20-21, 2024, at the Kigali Convention Centre, Kigali, Rwanda, was poised to be a landmark event in Rwanda's ongoing efforts to advance health research and policy. With the theme "Advancing Research for a Healthy Rwanda," this conference brought together a wide range of stakeholders from the health sector, including researchers, policymakers, healthcare professionals, academic institutions, and development partners. The event served as a platform for fostering collaboration, sharing knowledge, and exploring innovative approaches to addressing Rwanda's most pressing health challenges.

This conference was a pivotal occasion, including with the launch of two critical frameworks: the National Health Research Agenda (2024-2029) and the National Strategic Plan for Research Innovation and Data Science (2024-2029). Both documents represent comprehensive blueprints for prioritizing and guiding health research activities in Rwanda, ensuring that research efforts align with the country's health needs and strategic goals. The National Health Research Agenda focuses on leveraging research to address key health issues such as communicable and non-communicable diseases, maternal and child health, mental health, climate change and emergency health issues, antimicrobial resistance and others. Meanwhile, the Strategic Plan for Research Innovation and Data Science emphasizes the transformative potential of data-driven decision-making and technological innovations in healthcare.

One of the key goals of this conference was to create a connected ecosystem where research, policy, and practice intersect to improve healthcare delivery and outcomes. Rwanda, like many nations, faces a range of health challenges, including the burden of both communicable and non-communicable diseases and limited resources. The conference provided a platform for addressing the existing challenges through collaborative discussions and knowledge exchange among participants. In addition, the conference features special sessions focused on building human resource capacities in research and development (R&D), with a particular emphasis on fostering a culture of innovation. These sessions showcased strategies for enhancing the skills and expertise of health researchers in Rwanda, ensuring that the country is well-positioned to lead future advancements in healthcare.

This included the Three Minute Thesis (3MT) competition. This competition encouraged young researchers to showcase their innovative work, fostering a new generation of health researchers dedicated to addressing the unique health needs of Rwanda and the broader East African region.

Overall, the 2024 Health Research and Policy Conference was a vital step in strengthening Rwanda's health research ecosystem. It built on the success of previous achievements and commitments to promoting research as a key driver of healthcare improvement. The insights and innovations shared through this conference, reflected in the abstracts within this publication, serve as a valuable resource for advancing research, policy, and practice in Rwanda's health sector for years to come.

Potential Conflicts of Interest: No potential conflicts of interest disclosed by all authors. **Academic Integrity:** All authors confirm their substantial academic contributions to development of this manuscript as defined by the International Committee of Medical Journal Editors. **Originality:** All authors confirm this manuscript as an original piece of work, and confirm that has not been published elsewhere. **Review:** All authors allow this manuscript to be peer-reviewed by independent reviewers in a double-blind review process. © **Copyright:** The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY-NC-ND), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. **Publisher:** Rwanda Health Communication Centre, KG 302st., Kigali-Rwanda. Print ISSN: 2663 - 4651; Online ISSN: 2663 - 4653. **Website:** <https://rbc.gov.rw/publichealthbulletin/>

Very high epilepsy prevalence in rural Southern Rwanda: the underestimated burden of epilepsy in sub-Saharan Africa

Ieme Garrez^{1,*}, Dirk E Teuwen¹, Fidèle Sebera^{2,3}, Sylvestre Mutungirehe², Arlene Ndayisenga⁴, Delphine Kajeneza³, Georgette Umuhoza², Jeannine Kayirangwa⁵, Josiane Umwiringirwa², Peter Dedeken^{1,6}, Paul A.J.M. Boon¹

¹Department of Neurology, Ghent University Hospital, Ghent, Belgium

²Department of Neurology, Ndera Neuro-Psychiatric Teaching Hospital, Kigali, Rwanda

³Centre Hospitalier Universitaire Kigali, Rwanda

⁴King Faisal Hospital, Kigali, Rwanda

⁵Ruhengeri Referral Hospital, Musanze, Rwanda

⁶Heilig Hart Ziekenhuis, Lier, Belgium

ABSTRACT

Introduction: The burden of epilepsy in Rwanda is significant, particularly in rural areas, where recent studies have revealed very high prevalence rates. Our study assessed the prevalence of epilepsy across different geographical areas in Southern rural Rwanda

Methods: We undertook a prospective cross-sectional, door-to-door survey in a rural village in Southern Rwanda from June 2022 to April 2023. First, trained enumerators screened villagers for epilepsy using the validated Limoges epilepsy screening questionnaire. Second, neurologists examined persons who had screened positively to confirm epilepsy diagnosis.

Results: Enumerators screened 1,745 persons (54.4% female, mean age: 24 ± 19.3 years). Three hundred four persons (17.4%) screened positively. Epilepsy diagnosis was confirmed in 133 (52.6% female, mean age: 30 ± 18.2 years), and active epilepsy in 130 persons. Lifetime epilepsy prevalence was 76.2% (95%CI: 64.2-89.7%). The highest age-specific rate occurred in the 29-49 age group. No significant gender-specific differences were noted. The seizure onset occurred before 18 years of age in almost 60%. The diagnosis and treatment gaps were 79.4% and 92.2%, respectively. In 19.5% of the cases, only non-motor seizures occurred.

Conclusion: We report one of the highest epilepsy prevalence estimates documented, with almost 20% non-motor seizures, which are often underdiagnosed in rural Africa. In line with two previous Rwandan reports (2,3), we reiterate the burden of the disease in the country, with prevalence rates more than three times the estimated prevalence in SSA. Geographic variation in prevalence throughout Africa may result from differences in risk and etiological factors. Urgent case-control studies are underway to understand such differences to further clarify pathophysiological mechanisms and propose adapted epilepsy prevention policies.

***Corresponding author:**

Ieme Garrez
Ieme.Garrez@UGent.be
Tel: +32496138604

Department of Neurology, Ghent University Hospital, Ghent, Belgium

Received: September 4, 2024

Accepted: September 22, 2024

Published: September 30, 2024

Cite this article as: Garrez et al. Very high epilepsy prevalence in rural Southern Rwanda: the underestimated burden of epilepsy in sub-Saharan Africa. *Rw. Public Health Bul.* 2024, 5 (3): 8-8. <https://dx.doi.org/10.4314/rphb.v5i3.1>

Potential Conflicts of Interest: No potential conflicts of interest disclosed by all authors. **Academic Integrity:** All authors confirm their substantial academic contributions to development of this manuscript as defined by the International Committee of Medical Journal Editors. **Originality:** All authors confirm this manuscript as an original piece of work, and confirm that has not been published elsewhere. **Review:** All authors allow this manuscript to be peer-reviewed by independent reviewers in a double-blind review process. © **Copyright:** The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY-NC-ND), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. **Publisher:** Rwanda Health Communication Centre, KG 302st., Kigali-Rwanda. Print ISSN: 2663 - 4651; Online ISSN: 2663 - 4653. **Website:** <https://rbc.gov.rw/publichealthbulletin/>

Comparative Analysis of LAM Test and GeneXpert in Detecting Tuberculosis in Immunocompromised Patients

Evariste Ntezirizaza^{1,2,*}, Jeannine Uwimana^{1,2}, Chantal Mukandayishimiye¹, Fidele Habimana^{1,2}, Fidele Mahirane², Gilbert Uwizeyimana², Celestin Musabyumuremyi¹

¹Institute d'Enseignement Supérieur de Ruhengeli, Musanze, Rwanda

²Kigali University Teaching Hospital, Kigali, Rwanda

ABSTRACT

Introduction: Tuberculosis (TB) remains a significant global health concern, particularly among immunocompromised individuals, including those with HIV. Diagnosis of TB in such populations presents challenges due to atypical presentations and compromised immune responses. This study aimed to evaluate the diagnostic efficacy of LAM test, GeneXpert and TB culture, used in TB detection, especially, in immunocompromised patients at Kigali University Hospital (KUTH).

Methods: A cross-sectional study was conducted at KUTH on 74 samples of urine and sputum or another samples from suspected extrapulmonary sites. Urine samples were tested using the LAM test, GeneXpert, and culture, while sputum and extrapulmonary samples were examined using GeneXpert and culture only. The sensitivity of each method was assessed, with culture serving as the reference standard, and the data were analyzed using SPSS to compare the performance efficacy of LAM and GeneXpert in detecting TB in both HIV-positive and non-HIV patients.

Results: Among non-HIV patients, the LAM test demonstrated a sensitivity of 57% in detecting TB, while GeneXpert exhibited a slightly higher sensitivity of 75% among culture-positive cases. In HIV-positive patients, LAM exhibited superior sensitivity at 80%, compared to GeneXpert at 60% among the same participant pool.

Conclusion: This study highlights the varying diagnostic performance of LAM test and GeneXpert in detecting TB in immunocompromised patients, depending on HIV status. While LAM showed higher sensitivity in HIV-positive patients, GeneXpert performed marginally better in non-HIV cases. These findings underscore the importance of tailored diagnostic approaches based on patient immune status to optimize TB detection accuracy. Integration of LAM test and GeneXpert into TB diagnostic algorithms for immunocompromised populations could improve early detection and treatment initiation, thus reducing TB morbidity and mortality. Further research with larger cohorts is warranted to validate these findings and inform TB diagnostic strategies in resource-limited settings.

***Corresponding author:**

Evariste Ntezirizaza
Institute d'Enseignement Supérieur
de Ruhengeli, Musanze, Rwanda

ORCID: <https://orcid.org/0009-0001-7717-5971>

Received: September 4, 2024
Accepted: September 22, 2024
Published: September 30, 2024

Cite this article as: Ntezirizaza et al. Comparative Analysis of LAM Test and GeneXpert in Detecting Tuberculosis in Immunocompromised Patients: A Study at CHUK Hospital. *Rw. Public Health Bul.* 2024, 5 (3): 9-9. <https://dx.doi.org/10.4314/rphb.v5i3.1>

Potential Conflicts of Interest: No potential conflicts of interest disclosed by all authors. **Academic Integrity:** All authors confirm their substantial academic contributions to development of this manuscript as defined by the International Committee of Medical Journal Editors. **Originality:** All authors confirm this manuscript as an original piece of work, and confirm that has not been published elsewhere. **Review:** All authors allow this manuscript to be peer-reviewed by independent reviewers in a double-blind review process. © **Copyright:** The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY-NC-ND), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. **Publisher:** Rwanda Health Communication Centre, KG 302st., Kigali-Rwanda. Print ISSN: 2663 - 4651; Online ISSN: 2663 - 4653. **Website:** <https://rbc.gov.rw/publichealthbulletin/>

Knowledge, attitude, and practices towards tuberculosis prevention among HIV-positive patients at Kibagabaga District Hospital, Rwanda, 2021

Emile Sebera^{1,2,*}, Frank Kiiza^{1,3}, Janvier Irankunda¹, Consilie Mukarwego¹, Emile Twagirumukiza^{1,2}, Valens Bubanje^{1,2}

¹University of Rwanda, Kigali City, Rwanda

²Mount Kenya University, Kigali City, Rwanda

³Academy of Applied Sciences Higher School of Management and Administration, Opole, Poland

ABSTRACT

Introduction: Globally, an estimated 10.3 million people fell ill with TB in 2021, especially for people living with HIV. This study was conducted to assess the knowledge, attitudes, and practices (KAP) of TB prevention among HIV patients at Kibagabaga District Hospital (KDH). The study aimed to improve public health practices and the lives of people living with HIV by identifying knowledge gaps, tailoring prevention strategies, encouraging early TB detection, and informing policy development.

Methods: An analytical cross-sectional study was conducted at Kibagabaga District Hospital from November 2019 to February 2021. A sample size of 237 participants was calculated using Epi-info software and selected through a stratified simple random sampling method. Data collection employed self-administered questionnaires and analyzed using SPSS software version 23.

Results: Regarding the sociodemographic characteristics of the 237 HIV seropositive patients who participated in the study, the 39-48 age group represented the largest portion (35.4%), a majority were females by gender (60.3%), toward half (45.1%) being married, over half (52.5%) employed, nearly a half attained the primary education (51.1%). Upon the assessment of the knowledge level, attitude, and practice towards TB prevention, by aggregating the marks obtained from the respective questions, revealed a relative majority had a positive knowledge (86.9%), attitude (86.1%), and practices (91.6%) towards tuberculosis prevention.

Conclusion: While the study revealed positive overall knowledge, practices, and attitudes towards tuberculosis prevention among the HIV seropositive patients being followed up at Kibagabaga District Hospital, a potential knowledge gap regarding the specific cause of the disease was identified. This shows that improving health education efforts for HIV patients could benefit more. As far as comprehensive knowledge and ongoing positive behavior towards prevention are concerned, emphasis should be placed on understanding the causative agent of tuberculosis.

***Corresponding author:**

Emile Sebera
University of Rwanda, Bachelor of
Clinical Medicine and Community
Health, Kigali City, Rwanda

Email: emilesebera12@gmail.com

Received: September 4, 2024

Accepted: September 22, 2024

Published: September 30, 2024

Cite this article as: Sebera et al.
Knowledge, attitude, and practices
towards tuberculosis prevention
among HIV-positive patients at
Kibagabaga District Hospital,
Rwanda, 2021. *Rw. Public Health
Bul.* 2024. 5 (1): 10-10. [https://
dx.doi.org/10.4314/rphb.v5i3.3](https://dx.doi.org/10.4314/rphb.v5i3.3)

Potential Conflicts of Interest: No potential conflicts of interest disclosed by all authors. **Academic Integrity:** All authors confirm their substantial academic contributions to development of this manuscript as defined by the International Committee of Medical Journal Editors. **Originality:** All authors confirm this manuscript as an original piece of work, and confirm that has not been published elsewhere. **Review:** All authors allow this manuscript to be peer-reviewed by independent reviewers in a double-blind review process. © **Copyright:** The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY-NC-ND), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. **Publisher:** Rwanda Health Communication Centre, KG 302st., Kigali-Rwanda. Print ISSN: 2663 - 4651; Online ISSN: 2663 - 4653. **Website:** <https://rbc.gov.rw/publichealthbulletin/>

Classification of Long COVID patients using factors score Method: A case study of Rwanda

Rugwizangoga Eugene¹, Kayirangwa Marie Rose¹, Nsabiyeza Emile¹, Manirafasha Appolinaire², Turatsinze David^{3,*}, Rwagasore Edson⁴, Nizeyimana Pacifique⁵

¹Jhpiego-Rwanda, Kigali, Rwanda

²King Faisal Hospital, Kigali, Rwanda

³University teaching Hospital of Kigali, Kigali, Rwanda

⁴Rwanda Biomedical Centre, Kigali, Rwanda

⁵Adventist University of Central Africa, Kigali, Rwanda

ABSTRACT

Introduction: On a global scale, long COVID (LC) has not received the attention it deserves; there is limited knowledge on long COVID among healthcare providers and the public. In Rwanda, as of June 2023, 133,194 COVID-19 cases have been reported, and 131,647 have been recovered. It is known that the prevalence in the general population ranges from 10% to 30% of all cases of acute infection, and anyone with COVID-19 can get Long COVID-19. Understanding factors influencing long-term COVID-19 conditions can help clinicians improve the management of identified cases.

Methods: We have designed a tool in the form of a Likert scale questionnaire labeled Long Covid Screening Tool (LCST). The population targeted are patients visiting outpatient departments and non-communicable diseases clinics at the district hospital level from June 2023 to March 2024. Data was collected by doctors using REDCap. Through a summated Likert scale, we assign a total score to each observation; based on the total factor score and red flags symptoms, we classify patients into Grade I, Grade II, and Grade III.

Results: A total of 180 patients were included in the analysis. The most affected group was adults aged 30-55 years at 60%; females accounted for most cases at 65%; 98% were vaccinated with at least 2 shots. The most common comorbidities were hypertension 19% and diabetes mellitus 5%. The most reported symptoms were 64% fatigue and 57% memory loss. Based on the analysis of the score factors with grade I (0-13), grade II (14-26), and grade III (27-40), the majority, 83,3%, were classified as Grade I, 15,5%, are in Grade II, 1,2% in Grade III.

Conclusion: Classification of LC cases using the factor score method helped to know the severity. Results show that severe cases account for 1.2% of the cases, and those patients need to optimize their care. Further studies are required to classify the complexity of post-COVID-19 conditions. The sensitivity of health care providers and the integration of LC symptoms in routine clinical practice can help identify more cases and optimize their management.

*Corresponding author:
Dr. Turatsinze David
University teaching Hospital of
Kigali, Kigali, Rwanda

Email: dturatsinze2001@yahoo.fr

Received: September 4, 2024
Accepted: September 22, 2024
Published: September 30, 2024

Cite this article as: Turatsinze et al. Classification of Long COVID patients using factors score Method: a case study of Rwanda. *Rw. Public Health Bul.* 2024. 5 (3): 11-11. <https://dx.doi.org/10.4314/rphb.v5i3.1>

Potential Conflicts of Interest: No potential conflicts of interest disclosed by all authors. **Academic Integrity:** All authors confirm their substantial academic contributions to development of this manuscript as defined by the International Committee of Medical Journal Editors. **Originality:** All authors confirm this manuscript as an original piece of work, and confirm that has not been published elsewhere. **Review:** All authors allow this manuscript to be peer-reviewed by independent reviewers in a double-blind review process. © **Copyright:** The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY-NC-ND), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. **Publisher:** Rwanda Health Communication Centre, KG 302st., Kigali-Rwanda. Print ISSN: 2663 - 4651; Online ISSN: 2663 - 4653. **Website:** <https://rbc.gov.rw/publichealthbulletin/>

Prevalence of Undiagnosed Hypertension and its Associated Factors in Ndera Sector, Gasabo District of Rwanda: A Cross-Sectional Study

Akeem Ayodeji Okesina^{1,*}, Jean Claude Habineza², Richard Mbazumutima¹, Umurerwa Mignonne¹, Celine Mahirwe¹, Samuel Hakizimana¹, Francis Paul Wabwire¹

¹Department of Clinical Medicine and Community Health, School of Health Sciences, College of Medicine and Health Sciences, University of Rwanda, Kigali, Rwanda

²Rwandan Biomedical Center, NCDs Unit, Kigali, Rwanda

INTRODUCTION

Introduction: Hypertension remains a significant global health challenge, especially in low- and middle-income countries. It affects millions worldwide, with nearly half of the cases left undiagnosed. In Rwanda, a lack of adequate information and healthcare services impacts healthcare-seeking behaviors, contributing to undiagnosed hypertension in rural areas. Therefore, there exists a need to determine the incidence of undiagnosed hypertension and its associated factors.

Methods: A cross-sectional study of 393 adults was conducted in Ndera Sector, Gasabo District, through a multistage sampling technique. Data was collected using the WHO STEP-wise approach to non-communicable disease risk factor surveillance (STEPS) questionnaire; a physical examination was done for blood pressure and body mass index (BMI), after which the data collected were analyzed using SPSS.

Results: In Ndera Sector, 15% of the sample population had undiagnosed hypertension. Survey respondents had a mean age of 37 years (range 15 to 82), and 43% were male. Physical activity levels varied, with 41% reporting regular activity and 59% reporting a lack thereof. Significant factors associated with undiagnosed hypertension (p -value < 0.05) included age categories 45-54 years and ≥ 55 years, residing in Kibenga Cell, and having a BMI greater than 25.0 kg/m². Bivariate analysis revealed significant associations of undiagnosed hypertension with age (≥ 55), having a BMI greater than 30.0 kg/m² (obesity), and location of residence. The knowledge gap on hypertension also emerged as a significant factor, with those lacking awareness displaying a higher prevalence of undiagnosed hypertension. Sex, marital status, and other lifestyle habits showed no significant associations with undiagnosed hypertension.

Conclusion: A high prevalence of undiagnosed hypertension in the Ndera Sector was found to be associated with older age, high BMI, location of residence, and a lack of knowledge related to hypertension. This underscores the importance of health education initiatives, targeted healthcare initiatives in specific areas, tailored healthcare strategies for older age groups, and weight management initiatives to enhance awareness and early detection of hypertension.

*Corresponding author:
Dr Okesina Akeem Ayodeji
College of Medicine and Health
Sciences, University of Rwanda,
Email: akeemokesina@gmail.com

Received: September 4, 2024
Accepted: September 22, 2024
Published: September 30, 2024

Cite this article as: Okesina et al. Prevalence of Undiagnosed Hypertension and its Associated Factors in Ndera Sector, Gasabo District of Rwanda: A Cross-Sectional Study. *Rw. Public Health Bul.* 2024, 5 (3): 12-12. <https://dx.doi.org/10.4314/rphb.v5i3.1>

Potential Conflicts of Interest: No potential conflicts of interest disclosed by all authors. **Academic Integrity:** All authors confirm their substantial academic contributions to development of this manuscript as defined by the International Committee of Medical Journal Editors. **Originality:** All authors confirm this manuscript as an original piece of work, and confirm that has not been published elsewhere. **Review:** All authors allow this manuscript to be peer-reviewed by independent reviewers in a double-blind review process. © **Copyright:** The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY-NC-ND), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. **Publisher:** Rwanda Health Communication Centre, KG 302st., Kigali-Rwanda. Print ISSN: 2663 - 4651; Online ISSN: 2663 - 4653. **Website:** <https://rbc.gov.rw/publichealthbulletin/>

The prevalence of teeth replacement among edentulous patients attending the University of Rwanda Polyclinic

Habumugisha Jean Marie Vianney¹, Rutayisire Frank², Uwizeyimana Jean de Dieu³, Nyirahabimana Emeline⁴

¹*PaShen, Kigali, Rwanda*

²*Rwanda Military Hospital, Kigali, Rwanda*

³*Polyclinique la Medicale, Kigali, Rwanda*

⁴*Murunda District Hospital, Western Province, Rwanda*

ABSTRACT

Introduction: Despite the reduced rate of tooth loss, the need for prosthodontic treatment is still high. The most important reason to seek prosthetic replacement for missing teeth is to improve appearance, restoration of speech, mastication efficiency, confidence, and psychological well-being. Nevertheless, the prevalence of edentulous people remains high. However, Rwanda has a lack of dental laboratories, materials, and skilled personnel. But the few available are not utilized to the maximum since the report shows that the number of edentulous outweighs those that replaced their missing teeth, showing that much has to be done to increase awareness about aesthetics and function in terms of temporal mandibular joint pain and disfiguring of the face.

Objective of the study was to assess the prevalence of teeth replacement among the patients who attended the University of Rwanda Polyclinic.

Methods: In this retrospective descriptive study, data was collected using Excel and analyzed using SPSS version 25 from patients' medical files at the University of Rwanda Polyclinic.

Results: The 486 patients' medical files from January 2018 to January 2020 were used for data collection. There were 277 (57%) males and 209 (43%) females. Out of 486 patients, the prevalence of tooth loss was 267 (54.9%). The prevalence of tooth replacement was 44 (16.48%) out of 267 patients with tooth loss. Among these prostheses, 43 were acrylic removable partial dentures and 1 removable complete denture.

Conclusion: The findings have shown a high prevalence of tooth loss and a low prevalence of tooth replacement. Only a small number of participants have received prostheses. Universities and health institutions should reinforce the workforce through skilled laborers in dental clinics and laboratories and invest in improving oral health infrastructures.

***Corresponding author:**
Dr HABUMUGISHA Jean Marie Vianney
Email: habujmv@gmail.com

Received: September 4, 2024
Accepted: September 22, 2024
Published: September 30, 2024

Cite this article as: Habumugisha et al. The prevalence of teeth replacement among edentulous patients attending the University of Rwanda Poly-clinic. *Rw. Public Health Bul.* 2024. 5 (1): 13-13. <https://dx.doi.org/10.4314/rphb.v5i3.1>

Potential Conflicts of Interest: No potential conflicts of interest disclosed by all authors. **Academic Integrity:** All authors confirm their substantial academic contributions to development of this manuscript as defined by the International Committee of Medical Journal Editors. **Originality:** All authors confirm this manuscript as an original piece of work, and confirm that has not been published elsewhere. **Review:** All authors allow this manuscript to be peer-reviewed by independent reviewers in a double-blind review process. © **Copyright:** The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY-NC-ND), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. **Publisher:** Rwanda Health Communication Centre, KG 302st., Kigali-Rwanda. Print ISSN: 2663 - 4651; Online ISSN: 2663 - 4653. **Website:** <https://rbc.gov.rw/publichealthbulletin/>

Promoting the learning of Point-of-Care Ultrasound among undergraduate students through SonoGames: A Case Study from Rwanda

Ernest Aben Oumo^{1,*}, Beula Igiraneza¹, Axel Shimwa¹, Bruno Vargas², Chandler Villaverde^{1,3}, Miriam Frisch⁴, Onyinyechi Eke⁵, Kelsey Ripp¹

¹University of Global Health Equity, Rwanda

²Compañeros en Salud/Partners In Health, Mexico

³Inshuti Mu Buzima/Partners In Health, Rwanda

⁴Icahn School of Medicine at Mount Sinai, USA

⁵Massachusetts General Hospital, USA

ABSTRACT

Introduction: Point-of-care ultrasound (POCUS) has been shown to improve clinical diagnosis and medical decision-making. However, the incorporation of POCUS into medical education still requires robust curriculum development and assessment, especially in environments with low resources. Gamification of POCUS has been shown to be well received by learners, but its role in medical education on the African continent is still limited. In this study, we aimed to evaluate the effect of SonoGames on POCUS knowledge acquisition among medical students in Rwanda.

Methods: Three cohorts of University of Global Health Equity (UGHE) medical students in clinical rotations from 3rd year to 5th-year students participated in the SonoGames competition. The students had received prior education on POCUS at different points in their clinical training. They were invited to participate in “SonoGames”, a series of POCUS activities testing knowledge and skills acquisition. The students completed pre- and post-event surveys on POCUS knowledge, which consisted of image acquisition and image interpretation. Data were analyzed descriptively.

Results: 18 students participated in the SonoGames, including five 5th-year students, five 4th-year students, and eight 3rd-year students. 16/18 students (89%) completed the pre- and post-event surveys. In the pre-event survey, the mean grade on the ultrasound knowledge test was 75%. The mean grade on the post-event knowledge test was 86.5%. The students had very positive reviews of the session, with 14/16 students strongly agreeing that learning through the games improved their overall knowledge of POCUS, all students strongly agreeing that the games made them feel more enthusiastic about POCUS, and 15/16 students reporting that the games were a very effective educational experience. Additionally, 15/16 students strongly agreed that the games promoted team-building and student cooperation. In the survey results, most students requested ongoing SonoGames events at least annually.

Conclusion: Gamification of POCUS was very well received by students, who reported an increase in skills after the SonoGames event and demonstrated improved knowledge on pre- and post-testing. There is a need to continue to evaluate the curriculum so that medical students are equipped to practice clinically.

*Corresponding author:
Ernest Aben Oumo
University of Global Health Equity,
Rwanda
Email: ernest.oumo@student.ughe.org

Received: September 4, 2024
Accepted: September 22, 2024
Published: September 30, 2024

Cite this article as: Oumo et al. Promoting the learning of Point-of-Care Ultrasound among undergraduate students through SonoGames: A Case Study from Rwanda. *Rw. Public Health Bul.* 2024, 5 (3): 14-14. <https://dx.doi.org/10.4314/rphb.v5i3.1>

Potential Conflicts of Interest: No potential conflicts of interest disclosed by all authors. **Academic Integrity:** All authors confirm their substantial academic contributions to development of this manuscript as defined by the International Committee of Medical Journal Editors. **Originality:** All authors confirm this manuscript as an original piece of work, and confirm that has not been published elsewhere. **Review:** All authors allow this manuscript to be peer-reviewed by independent reviewers in a double-blind review process. © **Copyright:** The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY-NC-ND), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. **Publisher:** Rwanda Health Communication Centre, KG 302st., Kigali-Rwanda. Print ISSN: 2663 - 4651; Online ISSN: 2663 - 4653. **Website:** <https://rbc.gov.rw/publichealthbulletin/>

Mortality and sudden unexpected death in epilepsy in a cohort of 888 persons living with epilepsy in Rwanda

Cleme Garrez^{1,2}, Beni H. Uwacu³, Peter Dedeken^{1,2,4}, Fidèle Sebera^{3,5,6}, Josiane Umwiringirwa^{3,7}, Jeannine Kayirangwa⁸, Dirk E. Teuwen^{1,2}, Paul A.M.J. Boon²

¹Department of Neurology, Ghent University Hospital, Ghent, Belgium

²Brain, Ghent University Hospital, Ghent, Belgium

³Department of Neurology, Ndera Neuropsychiatric Teaching Hospital, Kigali, Rwanda

⁴Department of Neurology, Heilig Hart Ziekenhuis, Lier, Belgium

⁵Department of Neurology, King Faisal Hospital Rwanda, Kigali, Rwanda

⁶Department of Internal Medicine, Centre Hospitalier Universitaire Kigali, Kigali, Rwanda

⁷Division of Education, Training and Research, King Faisal Hospital Rwanda, Kigali, Rwanda

⁸Ruhengeri Referral Hospital, Musanze, Rwanda

ABSTRACT

Introduction: Mortality among people living with epilepsy is two- to three-fold higher than in the general population, with a 5.4 standardized mortality ratio in Sub-Saharan Africa. Sudden Unexpected Death in Epilepsy (SUDEP) is the most common cause of epilepsy-related death. We prospectively estimated mortality and SUDEP rates in Rwandan people living with epilepsy.

Methods: We conducted a longitudinal study at the tertiary Ndera Neuropsychiatric Teaching Hospital (Kigali) and at three primary healthcare centers in the Musanze District, Northern Province. People living with epilepsy, aged ≥ 15 years, were enrolled between February and December 2018. People living with epilepsy did not attend a close-out visit after 12 months were contacted. Upon report of a death, a neurologist traveled to meet family members and administered the WHO Verbal Autopsy Questionnaire to assess the cause of death.

Results: Out of a total of 888 people living with epilepsy, ten deaths were identified, yielding a mortality rate of 11.4/1000 person-years. Seven deaths had an identified cause, four of which were directly related to epilepsy, including death following head trauma, brain hematoma, prolonged seizure, and status epilepticus. No cause of death was identified in three cases, and these were interpreted as probable SUDEP, yielding a SUDEP rate of 3.4/1000 person-years. Mortality rates in tertiary and primary centers were comparable.

Conclusion: Mortality rates in people living with epilepsy are high and in line with previous reports from Rwanda. Improved treatment access and education of patients and families on death risks, including risk factors for SUDEP, should be provided to improve treatment compliance and better seizure control.

***Corresponding author:**

Cleme Garrez
Ghent University Hospital, Ghent,
Belgium
Email: ieme.garrez@ugent.be

Received: September 4, 2024
Accepted: September 22, 2024
Published: September 30, 2024

Cite this article as: Garrez et al. Mortality and sudden unexpected death in epilepsy in a cohort of 888 persons living with epilepsy in Rwanda. *Rw. Public Health Bul.* 2024. 5 (3): 15-15. <https://dx.doi.org/10.4314/rphb.v5i3.1>

Potential Conflicts of Interest: No potential conflicts of interest disclosed by all authors. **Academic Integrity:** All authors confirm their substantial academic contributions to development of this manuscript as defined by the International Committee of Medical Journal Editors. **Originality:** All authors confirm this manuscript as an original piece of work, and confirm that has not been published elsewhere. **Review:** All authors allow this manuscript to be peer-reviewed by independent reviewers in a double-blind review process. © **Copyright:** The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY-NC-ND), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. **Publisher:** Rwanda Health Communication Centre, KG 302st., Kigali-Rwanda. Print ISSN: 2663 - 4651; Online ISSN: 2663 - 4653. **Website:** <https://rbc.gov.rw/publichealthbulletin/>

Assessment of pseudomonas aeruginosa from trachea aspirates among Intensive Care Unit patients attending Rwanda Military Hospital

Lydivine Mpinganzima¹, Cedrick Michael Izere^{1,*}, Nemeyimana Patrick², Charles Muhinda³, Fabrice Uwumuremyi^{1,4}, Angeliqwe Usengimana⁴, Berchmas Mutijima^{1,4}

¹Ines Ruhengeri, Musanze, Rwanda

²University of Rwanda, Kigali, Rwanda

³Rwanda Military referral and teaching hospital, Kigali, Rwanda

⁴Legacy clinics and diagnostics ltd, Kigali, Rwanda

ABSTRACT

Introduction: Pseudomonas aeruginosa (PA) is a non-fermenting Gram-negative bacillus that frequently causes nosocomial infections, particularly in the intensive care unit (ICU), contributing to substantial mortality. They are responsible for ventilator-associated pneumonia and hospital-acquired pneumonia, where they account for 24% of ICU respiratory infections. This research aimed to determine the rate of Pseudomonas aeruginosa among tracheal aspirates and assess its susceptibility towards different antibiotics at Rwanda Military Hospital, which could contribute to the infectious control team to design appropriate strategies for prevention of Pseudomonas aeruginosa-associated respiratory infection and antibiotic resistance stewardship.

Method: This is a retrospective study where the data of the patients suspected of respiratory tract infection hospitalized in the intensive care unit from January 2021 to June 2023 were collected and analyzed. The collected data were presented using Microsoft Excel 2016, and data were analyzed using SPSS Version 16 statistical software.

Results: Out of 414 data collected, 181 (43.7%) of samples reported growth of at least one organism (isolate); among them 89 (49.1%) were Pseudomonas aeruginosa, and 92 (50.8%) were positive for other species. Isolated Pseudomonas aeruginosa was sensitive to only PB (92.6%), totally resistant to CAZ (98.5%), and half of first-line antibiotics used above 80%.

Conclusion: The rate of Pseudomonas aeruginosa isolates was found to be high in intensive care units, potentially being a threat as it is known to be a strong hospital-acquired infection. The same applies to antibiotic susceptibility tests where resistance was found high. There is a need to provide measures to minimize the spread of resistance of Pseudomonas aeruginosa, which is observed to develop towards antibiotics more quickly than in most bacteria, and improvement in hospitals towards the use of aseptic techniques while performing procedures and working in infected environments also disinfection while using mechanical ventilation.

***Corresponding author:**

Dr. Cedrick Michael IZERE
Department biomedical laboratory sciences, Ines Ruhengeri, Musanze, Rwanda
Email: icedrick@ines.ac.rw

Received: September 4, 2024

Accepted: September 22, 2024

Published: September 30, 2024

Cite this article as: Mpinganzima et al. Assessment of pseudomonas aeruginosa from trachea aspirates among intensive care unit patients attending Rwanda military hospital. *Rw. Public Health Bul.* 2024. 5 (3): 16-16. <https://dx.doi.org/10.4314/rphb.v5i3.1>

Potential Conflicts of Interest: No potential conflicts of interest disclosed by all authors. **Academic Integrity:** All authors confirm their substantial academic contributions to development of this manuscript as defined by the International Committee of Medical Journal Editors. **Originality:** All authors confirm this manuscript as an original piece of work, and confirm that has not been published elsewhere. **Review:** All authors allow this manuscript to be peer-reviewed by independent reviewers in a double-blind review process. © **Copyright:** The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY-NC-ND), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. **Publisher:** Rwanda Health Communication Centre, KG 302st., Kigali-Rwanda. Print ISSN: 2663 - 4651; Online ISSN: 2663 - 4653. **Website:** <https://rbc.gov.rw/publichealthbulletin/>

Assessment of Intestinal Parasites and Risk Factors Among HIV Patients Attending Ntaruka Health Center, Burera District, Rwanda

Musabyumuremyi Celestin^{1,*}, Mukashema Hyacinthe², Nyiramahirwe Denyse²

¹University of Rwanda, College of Medicine and Health Sciences, Kigali, Rwanda

²Institut d'Enseignement Supérieur de Ruhengeri (INES), Musanze, Rwanda

ABSTRACT

Introduction: HIV infection is still a public health problem worldwide and co-infections with other infectious agents, including intestinal parasites, are of particular concern, mainly in developing countries such as Rwanda. Therefore, understanding the epidemiology of intestinal parasitic infection is essential for the effective management of HIV infection. The aim of this study was to assess the burden of intestinal parasites and risk factors among HIV patients attending Ntaruka Health Center located in Burera District and this helps to take appropriate measures in the management of people living with HIV.

Methods: This was a cross-section study conducted from April to August 2023, where a total sample of 80 HIV-positive patients participated in the study. A questionnaire was designed to assess the risk factors. Stool samples were examined microscopically to assess the diagnostic forms of different parasites and negative sample results were concentrated using Formal – Ether technique. Blood samples were collected to analyze HIV viral Load and CD4. All data were presented and analyzed in SPSS version 22, and the Chi-square test was used to analyze the results.

Results: Among 80 participants, 54 individuals were infected with intestinal parasites, with a prevalence of (67.5%). The most prevalent intestinal parasites were *Ascaris lumbricoides* 25%, *Trichuris trichiura* (17.5%), *Giardia lamblia* 13.75%; *Entamoeba histolytica* to 11.25%. In relation to gender, females were found to be infected than males at 63.8% and 36.2%, respectively. The illiterate individual was 53.75% compared to 46.25 % for literate. According to age groups, the highest predominant age group was the group above 50 years old (36.25%), the following were the age groups ranging from 41-50 years old (28.75%), 31-40 (21.25%) and the last age group was from 15-30 (13.75%). The following risk factors were statistically associated with parasitic infections: using water from lakes and rivers as domestic sources, eating unwashed vegetables and fruits, defecating in bushes near farms during farming, and using human feces as fertilizer. This study had no significant correlation between intestinal parasites with the number of CD4+ T cells and HIV Viral Load.

Conclusion: HIV patients should be screened routinely for intestinal parasites, and more research are needed countrywide in order to assess the magnitude of the problem.

***Corresponding author:**

Celestin MUSABYUMUREMYI
University of Rwanda, College
of Medicine and Health Sciences,
Kigali, Rwanda
Email: cemusabu@gmail.com

Received: September 4, 2024

Accepted: September 22, 2024

Published: September 30, 2024

Cite this article as:
Musabyumuremyi et al.
Assessment of Intestinal Parasites
and Risk Factors Among HIV
Patients Attending Ntaruka Health
Center, Burera District, Rwanda.
Rw. Public Health Bul. 2024, 5 (3):
17-17. [https://dx.doi.org/10.4314/
rphb.v5i3.1](https://dx.doi.org/10.4314/rphb.v5i3.1)

Potential Conflicts of Interest: No potential conflicts of interest disclosed by all authors. **Academic Integrity:** All authors confirm their substantial academic contributions to development of this manuscript as defined by the International Committee of Medical Journal Editors. **Originality:** All authors confirm this manuscript as an original piece of work, and confirm that has not been published elsewhere. **Review:** All authors allow this manuscript to be peer-reviewed by independent reviewers in a double-blind review process. © **Copyright:** The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY-NC-ND), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. **Publisher:** Rwanda Health Communication Centre, KG 302st., Kigali-Rwanda. Print ISSN: 2663 - 4651; Online ISSN: 2663 - 4653. **Website:** <https://rbc.gov.rw/publichealthbulletin/>

Evaluating the impact of Community-Based Socioterapy on social dignity and mental health of different social groups in post-genocide Rwanda: A clustered Randomised Controlled Trial

Sandra Schwarz^{1,*}, Japhet Niyonsenga¹, Epaphrodite Nsabimana¹, Angela Jansen², Chantal Marie Ingabire³, Stefan Jansen¹

¹*Mental Health & Behaviour Research Group, University of Rwanda, Kigali, Rwanda*

²*Institute for Community-Based Socioterapy (ICBS), Kigali, Rwanda*

³*Center for Community-Based Socioterapy (CBS Rwanda), Kigali, Rwanda*

ABSTRACT

Introduction: Community-Based Socioterapy (CBS) is a psychosocial community intervention implemented in post-genocide Rwanda since 2005. It aims at enhancing social dignity (an innovative construct in mental health research) and mental health within Rwandan communities. This study marks the first clustered Randomised Controlled Trial (RCT) to evaluate and compare the intervention's effects on social dignity and mental health among participating Genocide Survivors, Ex-Prisoners, and Others (people who were neither directly targeted nor perpetrators).

Methods: The study included 80 groups of 10-15 individuals from ten Rwandan districts, giving us a total of 833 participants. Half of these groups were randomly selected as intervention groups, while the other half were waitlisted. The intervention groups completed the CBS program over 15 weeks. Data was collected at baseline, endline, and 4.5 months post-intervention. Social dignity was evaluated through a newly developed Social Dignity Scale (SDS). Mental health was assessed through depressive symptoms (PHQ-9), PTSD symptoms (PCL-5), and psychological well-being (WHO-5). Statistical analyses were conducted in R using ANOVA and LMM.

Results: The analyses revealed that baseline differences between Survivors, Ex-Prisoners and Others in social dignity and mental health were largely attributable to covariates such as gender, relationship status, and trauma exposure. The intervention significantly improved both social dignity and mental health (Pseudo-R²: 0.04 approx.). No significant differences concerning the intervention's positive impact were found between Genocide Survivors, Ex-Prisoners, and Others.

Conclusion: Our results suggest that CBS effectively enhances social dignity and mental health regardless of participant's social group. Future research should explore the rather small effect size observed in this study.

*Corresponding author:

Sandra Schwarz
Mental Health & Behaviour
Research Group, University of
Rwanda, Kigali, Rwanda

Email: sandra.m.schwarz@gmail.com

Received: September 4, 2024
Accepted: September 22, 2024
Published: September 30, 2024

Cite this article as: Schwarz et al. Evaluating the impact of Community-Based Socioterapy on social dignity and mental health of different social groups in post-genocide Rwanda: A clustered Randomised Controlled Trial. *Rw. Public Health Bul.* 2024, 5 (3): 18-18. <https://dx.doi.org/10.4314/rphb.v5i3.1>

Potential Conflicts of Interest: No potential conflicts of interest disclosed by all authors. **Academic Integrity:** All authors confirm their substantial academic contributions to development of this manuscript as defined by the International Committee of Medical Journal Editors. **Originality:** All authors confirm this manuscript as an original piece of work, and confirm that has not been published elsewhere. **Review:** All authors allow this manuscript to be peer-reviewed by independent reviewers in a double-blind review process. © **Copyright:** The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY-NC-ND), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. **Publisher:** Rwanda Health Communication Centre, KG 302st., Kigali-Rwanda. Print ISSN: 2663 - 4651; Online ISSN: 2663 - 4653. **Website:** <https://rbc.gov.rw/publichealthbulletin/>

Using Natural Language Processing to Evaluate Home-based care text exchange between patients and care provider during COVID-19 Pandemic in Rwanda: A Model to Support rapid Pandemic Decision Making

Hassan Mugabo¹, Matthew Manson¹, Muhammed Semakula², Hyeju Jang², Guiseppe Carenini¹, Richard T Lester¹

¹Research Innovation and Data Science Divison (RIDS), Rwanda Biomedical Centre (RBC), Kigali, Rwanda

²Rwanda Ministry of Health, Kigali, Rwanda

ABSTRACT

***Corresponding author:**

Hassan Mugabo
Rwanda Biomedical Centre, Kigali,
Rwanda

Email: mugaboden@gmail.com

Received: September 4, 2024

Accepted: September 22, 2024

Published: September 30, 2024

Cite this article as: Mugabo et al. Using Natural Language Processing to Evaluate Home-based care text exchange between patients and care provider during COVID-19 Pandemic in Rwanda: A Model to Support rapid Pandemic Decision Making. *Rw. Public Health Bul.* 2024. 5 (3): 19-19. <https://dx.doi.org/10.4314/rphb.v5i3.1>

Introduction: Isolation of cases is a cornerstone of public health measures to contain the spread of pathogens. The Rwandan government deployed a multi-language digital health tool that facilitated daily text message communication with patients diagnosed with SARS-CoV-2 cases and their contacts as part of national COVID-19 home-based care program. We evaluated the texting patterns and communication topics to gain insights into patient experiences in these circumstances.

Methods: We extracted data on all COVID-19 patients (cases and contacts) enrolled in the WelTel text messaging program in Rwanda between March 18, 2020 and March 31, 2022 and linked demographic and clinical data from the national COVID-19 registry. We examined texting patterns and message content from users. A portion of the total conversation corpus was translated into English and labeled with pre-defined medical and social topics of interest identified by medical experts.

Several NLP topic classification models were trained and tested using the labelled untranslated or English-translated conversations, and compared for performance using F1-scores. The best performing models were applied to the unlabeled and untranslated conversations to label and classify the complete corpus. Topic frequencies were compared by demographic and clinical factors.

Results: In total, 33,081 patients, with mean age of 33.9 years old (0-100 years), and (44% female) including 30,398 cases and 2,683 contacts were registered in WelTel over 2 years of COVID-19 waves. This generated 12,119 interactive text conversations between patients and healthcare providers. Transformer models tended to yield better F1 performance scores than classical models when classifying English translated data, but classical machine models performed better when applied to the untranslated (mostly Kinyarwanda) corpus. The most frequent topic discussed was reporting whether symptoms were present. Discussions about prevention (e.g., transmission risk or vaccines), management (e.g. treatment of fever), social stressors (e.g. hunger or school attendance), and culture (e.g., religious references) also occurred. Targeted advice and education were frequently provided to patients who used the service in real-time.

Conclusion: Open language interactive text message-based systems can remotely support cases and contacts management in pandemics. NLP is a powerful computational approach that, if improved, could provide timely insights into the medical and social stressors that affect isolated patients.

Potential Conflicts of Interest: No potential conflicts of interest disclosed by all authors. **Academic Integrity:** All authors confirm their substantial academic contributions to development of this manuscript as defined by the International Committee of Medical Journal Editors. **Originality:** All authors confirm this manuscript as an original piece of work, and confirm that has not been published elsewhere. **Review:** All authors allow this manuscript to be peer-reviewed by independent reviewers in a double-blind review process. © **Copyright:** The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY-NC-ND), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. **Publisher:** Rwanda Health Communication Centre, KG 302st., Kigali-Rwanda. Print ISSN: 2663 - 4651; Online ISSN: 2663 - 4653. **Website:** <https://rbc.gov.rw/publichealthbulletin/>

Evaluation of risk factors for poorly controlled diabetes through glycated hemoglobin among diabetic patients attending Rwamagana level two teaching hospital, July 2023

Jean Paul Batibuka^{1,*}, Celestin Musabyumuremyi², Consolée Uwamahoro²

¹Rwamagana Referral Hospital, Rwamagana, Rwanda

²Institut d'Enseignement Supérieur de Ruhengeri (INES), Musanze, Rwanda

ABSTRACT

Introduction: The prevalence of diabetes in Rwanda has reached 3% of the total population, and other countries such as South Africa, Uganda, and Tanzania are reaching 8.39%, 4.42%, and 7.95%, respectively. For these patients, clinical management is essential to prevent morbidity and mortality. Glycated hemoglobin plays a vital role in diabetes management. An ideal value of <7% demonstrates control over the disease and enables prevention of its complications. This study in Rwamagana Hospital aims to understand the factors associated with poor prognosis of diabetes and recommend appropriate measures.

Methods: This study was cross-sectional prospective. Laboratory analyses were done to test HbA1c in blood, and medical files were reviewed. The study population was the clients with type 2 diabetes mellitus (236) received in July 2023 who accepted to participate voluntarily in the study, with a sample size of 149 participants, who were selected randomly. The difference in proportions by various clinical factors such as weight, sex, and medications were tested for their association with the outcome of having poorly controlled diabetes (HbA1c $\geq 7\%$). The chi-square test was used to determine the association between variables. The significance was defined at p-value < 0.05.

Results: Among those with poorly controlled diabetes, 36.2% were males and 63.8% were females. For body mass index, 49.7% were normal weight and 50.3% were overweight/obese. There was a significant association between sex and diabetes control, with female patients being more likely to have poorly controlled diabetes (P-value < 0.032). Being overweight/obese was significantly associated with having poor control of diabetes (P-value = 0.039).

Conclusion: People with diabetes under follow-up at Rwamagana level two teaching hospital have increased HbA1c levels, which is linked with gender and body mass index, which require different stakeholders to mobilize the population to do physical exercises and consume a balanced diet to avoid obesity.

***Corresponding author:**

Jean Paul Batibuka
Rwamagana Referral Hospital,
Rwamagana, Rwanda

Email: batibuka2003@gmail.com

Received: September 4, 2024

Accepted: September 22, 2024

Published: September 30, 2024

Cite this article as: Batibuka et al. Evaluation of risk factors for poorly controlled diabetes through glycated hemoglobin among diabetic patients attending Rwamagana level two teaching hospital, July 2023. *Rw. Public Health Bul.* 2024, 5 (3): 20-20. <https://dx.doi.org/10.4314/rphb.v5i3.1>

Potential Conflicts of Interest: No potential conflicts of interest disclosed by all authors. **Academic Integrity:** All authors confirm their substantial academic contributions to development of this manuscript as defined by the International Committee of Medical Journal Editors. **Originality:** All authors confirm this manuscript as an original piece of work, and confirm that has not been published elsewhere. **Review:** All authors allow this manuscript to be peer-reviewed by independent reviewers in a double-blind review process. © **Copyright:** The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY-NC-ND), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. **Publisher:** Rwanda Health Communication Centre, KG 302st., Kigali-Rwanda. Print ISSN: 2663 - 4651; Online ISSN: 2663 - 4653. **Website:** <https://rbc.gov.rw/publichealthbulletin/>

Actors and Channels Preferred by Youth for Accessing Mental Health-Related Services in Rwanda

Elvis Nininahazwe^{1,*}, Darius Gishoma², and Stefan Jansen¹

¹College of Medicine and Health Sciences, University of Rwanda, Kigali, Rwanda

²Rwanda Biomedical Centre, Mental health division, Kigali, Rwanda

ABSTRACT

Introduction: Following the 1994 genocide against Tutsi in Rwanda, government and international organizations implemented mental health services to support those suffering from the detrimental consequences. Despite the availability of these services, only 25% of individuals with reported mental disorders seek professional care, and the situation may be even worse for Rwandan youth based on anecdotal evidence. This study aimed to investigate youth's perspectives on access to mental health care; to explore 'actors' and 'channels' utilized by Rwandan youth for accessing mental health information and dealing with mental health-related issues.

Methods: This exploratory qualitative study involved 29 semi-structured interviews and two focus group discussions with youth who were purposively selected from both rural and urban areas of Rwanda. Data analysis employed thematic analysis with both deductive and inductive approaches to address the study objectives.

Results: This study revealed disparities in identifying factors influencing mental health among youth. Those from underprivileged families attributed mental health issues to spiritual causes, while those from privileged families cited family conflicts and societal pressures. Factors such as lived experiences, and socioeconomic status significantly shaped youth engagement with mental health information and services. Barriers, including lack of trust in actors and channels, lack of awareness, financial constraints, and stigma, hindered youth from engaging with mental health. According to the participants, addressing these barriers necessitates targeted awareness campaigns and youth-led initiatives to enhance mental health literacy and access to support services. For those seeking mental health-related information and/or support, friends, family members, nurses, and doctors were commonly used actors. Social media platforms, the internet, radio, peer discussions, and therapy were the most utilized channels.

Conclusion: Our findings underline the major challenges faced by Rwandan youth in accessing mental health services and information and identify the actors and channels used. The study contributes to the growing body of knowledge on mental health service access in Rwanda. It also offers valuable insights for policymakers, organizations, and individuals seeking to improve mental health services for Rwandan youth. By addressing the identified barriers, they can foster a more supportive environment for youth to seek and receive mental health support.

*Corresponding author:

Elvis Nininahazwe
College of Medicine and Health
Sciences, University of Rwanda,
Kigali, Rwanda

Email: elnininahazwe@gmail.com

Received: September 4, 2024

Accepted: September 22, 2024

Published: September 30, 2024

Cite this article as: Nininahazwe et al. Actors and Channels Preferred by Youth for Accessing Mental Health-Related Services in Rwanda. *Rw. Public Health Bul.* 2024, 5 (3): 21-21. <https://dx.doi.org/10.4314/rphb.v5i3.1>

Potential Conflicts of Interest: No potential conflicts of interest disclosed by all authors. **Academic Integrity:** All authors confirm their substantial academic contributions to development of this manuscript as defined by the International Committee of Medical Journal Editors. **Originality:** All authors confirm this manuscript as an original piece of work, and confirm that has not been published elsewhere. **Review:** All authors allow this manuscript to be peer-reviewed by independent reviewers in a double-blind review process. © **Copyright:** The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY-NC-ND), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. **Publisher:** Rwanda Health Communication Centre, KG 302st., Kigali-Rwanda. Print ISSN: 2663 - 4651; Online ISSN: 2663 - 4653. **Website:** <https://rbc.gov.rw/publichealthbulletin/>